

MY 2022 (RY 2023) HEDIS® Rate Chart

| Measure | Health Plan | | | | | | | MY 2022 HC Weighted Average | MY 2021 HC Weighted Average | MY 2020 HC Weighted Average |
|--|----------------------------------|-----------------------|-----------------------|---------------------------------|----------------|-------------------|--------------|-----------------------------|-----------------------------|-----------------------------|
| | AmeriHealth Caritas Pennsylvania | Geisinger Health Plan | Health Partners Plans | Highmark Wholecare ¹ | Keystone First | United Healthcare | UPMC for You | | | |
| Domain: Effectiveness of Care | | | | | | | | | | |
| BMI Percentile Documentation, Ages 3 to 11 Years ² | 80.57% | 82.87% | 88.02% | 86.69% | 85.37% | 88.56% | 79.76% | 83.61% | 83.90% | 80.77% |
| BMI Percentile Documentation, Ages 12 to 17 Years ² | 68.46% | 80.95% | 80.60% | 85.81% | 86.21% | 85.71% | 80.53% | 80.84% | 81.55% | 76.52% |
| BMI Percentile Documentation, Total ² | 75.41% | 82.16% | 85.37% | 86.37% | 85.67% | 87.59% | 80.00% | 82.48% | 83.02% | 79.30% |
| Counseling for Nutrition, Ages 3 to 11 Years ² | 76.57% | 70.83% | 78.93% | 79.47% | 76.10% | 82.66% | 72.47% | 75.71% | 78.16% | 74.75% |
| Counseling for Nutrition, Ages 12 to 17 Years ² | 64.62% | 74.60% | 80.60% | 72.97% | 68.10% | 85.00% | 70.80% | 71.52% | 74.20% | 71.56% |
| Counseling for Nutrition, Total ² | 71.48% | 72.22% | 79.52% | 77.13% | 73.21% | 83.45% | 71.94% | 74.10% | 76.62% | 73.64% |
| Counseling for Physical Activity, Ages 3 to 11 Years ² | 71.43% | 67.59% | 65.70% | 74.90% | 69.76% | 77.49% | 70.04% | 70.30% | 72.97% | 68.07% |
| Counseling for Physical Activity, Ages 12 to 17 Years ² | 63.85% | 75.40% | 80.60% | 69.59% | 72.41% | 83.57% | 71.68% | 72.16% | 74.19% | 69.99% |
| Counseling for Physical Activity, Total ² | 68.20% | 70.47% | 71.01% | 72.99% | 70.72% | 79.56% | 70.56% | 70.88% | 73.45% | 68.78% |
| Childhood Immunization Status, DtaP/DT | 75.67% | 74.70% | 72.26% | 75.67% | 73.24% | 73.24% | 70.32% | 73.31% | 74.49% | 76.80% |
| Childhood Immunization Status, IPV | 88.08% | 91.73% | 85.64% | 89.05% | 86.37% | 87.59% | 87.35% | 87.71% | 89.61% | 91.24% |
| Childhood Immunization Status, MMR | 87.59% | 86.62% | 87.35% | 87.35% | 85.89% | 85.16% | 87.35% | 86.82% | 86.76% | 89.07% |
| Childhood Immunization Status, HiB | 86.37% | 86.62% | 85.40% | 87.35% | 86.62% | 85.40% | 85.89% | 86.28% | 87.65% | 89.07% |
| Childhood Immunization Status, Hepatitis B | 89.05% | 92.21% | 89.29% | 91.24% | 86.37% | 90.27% | 89.54% | 89.30% | 91.24% | 92.21% |
| Childhood Immunization Status, VZV | 85.64% | 86.86% | 87.35% | 87.35% | 86.37% | 85.16% | 86.62% | 86.50% | 86.56% | 88.49% |
| Childhood Immunization Status, Pneumococcal Conjugate | 76.40% | 78.83% | 74.45% | 77.86% | 73.97% | 74.70% | 73.97% | 75.41% | 76.64% | 79.12% |
| Childhood Immunization Status, Hepatitis A | 81.75% | 84.18% | 85.89% | 85.16% | 82.73% | 81.51% | 83.70% | 83.51% | 83.76% | 85.27% |
| Childhood Immunization Status, Rotavirus | 68.61% | 75.18% | 68.37% | 70.80% | 73.24% | 73.48% | 71.53% | 71.53% | 72.23% | 73.65% |
| Childhood Immunization Status, Influenza | 40.39% | 42.58% | 52.07% | 43.31% | 54.74% | 50.12% | 37.71% | 45.64% | 50.84% | 53.35% |
| Childhood Immunization Status, Combo 3 | 68.61% | 69.83% | 67.88% | 70.56% | 68.86% | 69.34% | 63.75% | 67.97% | 69.32% | 72.15% |
| Childhood Immunization Status, Combo 7 | 58.39% | 62.29% | 58.64% | 59.61% | 62.04% | 61.56% | 54.01% | 59.06% | 59.05% | 62.16% |
| Childhood Immunization Status, Combo 10 | 32.36% | 34.06% | 41.85% | 34.31% | 45.26% | 40.88% | 27.74% | 36.36% | 39.23% | 41.83% |
| Immunizations for Adolescents, Combo 1 | 86.86% | 86.86% | 86.13% | 89.29% | 85.92% | 83.45% | 88.81% | 86.98% | 84.78% | 87.62% |
| Lead Screening in Children ³ | 80.47% | 86.62% | 74.19% | 85.40% | 81.79% | 79.45% | 83.94% | 81.92% | 81.60% | 83.22% |
| Breast Cancer Screening, Ages 52 to 74 Years ² | 58.09% | 57.39% | 54.51% | 50.68% | 53.41% | 48.47% | 56.40% | 55.06% | 50.90% | 53.16% |
| Cervical Cancer Screening, Ages 21 to 64 Years ² | 63.27% | 62.16% | 57.32% | 60.34% | 55.59% | 51.09% | 57.11% | 58.45% | 60.51% | 61.11% |
| Colorectal Cancer Screening (Total) ⁴ | 39.99% | 42.09% | 33.57% | 38.86% | 33.84% | 27.10% | 44.67% | 38.96% | | |
| Non-Recommended Cervical Cancer Screening in Adolescent Females, Ages 16 to 20 Years | 0.20% | 0.86% | 0.03% | 0.15% | 0.10% | 0.08% | 0.27% | 0.24% | 0.33% | 0.39% |
| Chlamydia Screening in Women, Ages 16 to 20 Years | 46.79% | 45.53% | 64.57% | 49.90% | 63.88% | 63.76% | 46.94% | 53.43% | 53.17% | 53.67% |
| Chlamydia Screening in Women, Ages 21 to 24 Years | 56.57% | 55.73% | 68.37% | 60.04% | 69.89% | 68.47% | 56.13% | 61.60% | 61.36% | 61.01% |
| Appropriate Testing for Pharyngitis, Ages 3 to 17 Years ² | 68.28% | 68.38% | 80.54% | 77.35% | 81.99% | 79.11% | 79.52% | 75.73% | 73.72% | 82.07% |
| Appropriate Testing for Pharyngitis, Ages 18 to 64 Years ² | 51.46% | 55.37% | 39.23% | 60.28% | 38.95% | 59.91% | 59.02% | 53.44% | 51.89% | 59.60% |
| Appropriate Testing for Pharyngitis, Ages 65 Years and Older ² | NA | NA | NA | NA | NA | NA | 33.33% | 33.33% | NA | NA |
| Appropriate Testing for Pharyngitis, Total ² | 60.99% | 62.90% | 58.24% | 69.94% | 60.45% | 69.63% | 70.34% | 65.55% | 62.14% | 74.25% |

MY 2022 (RY 2023) HEDIS® Rate Chart

| Measure | Health Plan | | | | | | | MY 2022 HC Weighted Average | MY 2021 HC Weighted Average | MY 2020 HC Weighted Average |
|---|----------------------------------|-----------------------|-----------------------|---------------------------------|----------------|-------------------|--------------|-----------------------------|-----------------------------|-----------------------------|
| | AmeriHealth Caritas Pennsylvania | Geisinger Health Plan | Health Partners Plans | Highmark Wholecare ¹ | Keystone First | United Healthcare | UPMC for You | | | |
| Appropriate Treatment for Upper Respiratory Infection, Ages 3m to 17 Years | 94.41% | 92.19% | 97.75% | 95.58% | 97.61% | 96.01% | 94.33% | 95.08% | 95.66% | 94.21% |
| Appropriate Treatment for Upper Respiratory Infection, Ages 18 to 64 Years | 86.78% | 83.80% | 82.41% | 88.12% | 78.71% | 86.33% | 84.67% | 84.94% | 85.53% | 81.97% |
| Appropriate Treatment for Upper Respiratory Infection, Ages 65 Years and Older | 83.87% | NA | 64.15% | NA | 73.97% | NA | 72.19% | 72.32% | 89.36% | 77.77% |
| Appropriate Treatment for Upper Respiratory Infection, Total | 92.40% | 89.92% | 94.39% | 93.66% | 94.37% | 93.34% | 91.44% | 92.45% | 92.38% | 90.89% |
| Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis, Ages 3m to 17 Years | 74.38% | 69.83% | 88.02% | 77.38% | 85.13% | 83.49% | 76.80% | 78.16% | 76.09% | 73.79% |
| Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis, Ages 18 to 64 Years | 50.08% | 52.27% | 61.19% | 52.42% | 51.11% | 48.11% | 46.80% | 50.49% | 49.49% | 46.26% |
| Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis, Ages 65 Years and Older | NA | NA | NA | NA | NA | NA | 36.27% | 36.27% | NA | NA |
| Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis, Total | 65.62% | 61.78% | 76.14% | 68.19% | 74.69% | 68.96% | 62.09% | 66.72% | 60.03% | 60.67% |
| Use of Spirometry Testing in the Assessment and Diagnosis of Chronic Obstructive Pulmonary Disease (COPD) | 25.49% | 27.59% | 18.69% | 25.08% | 22.68% | 22.02% | 22.94% | 23.42% | 24.17% | 26.86% |
| Pharmacotherapy Management of COPD Exacerbation (Systemic Corticosteroids) | 82.43% | 78.33% | 71.90% | 79.52% | 75.77% | 71.29% | 80.52% | 78.32% | 77.97% | 77.23% |
| Pharmacotherapy Management of COPD Exacerbation (Bronchodilators) | 90.60% | 86.69% | 88.93% | 87.01% | 89.90% | 83.13% | 88.74% | 88.33% | 87.20% | 87.29% |
| Statin Therapy for Patients With Cardiovascular Disease Received Statin Therapy (Total) ² | 85.00% | 85.91% | 83.99% | 83.93% | 82.42% | 80.72% | 84.59% | 84.24% | 84.24% | 83.51% |
| Statin Therapy for Patients With Cardiovascular Disease Statin Adherence 80% (Total) ² | 78.41% | 78.64% | 73.63% | 75.45% | 78.55% | 71.34% | 80.65% | 78.41% | 75.57% | 76.32% |
| Statin Therapy for Patients With Diabetes Received Statin Therapy (Total) ² | 67.99% | 66.30% | 72.28% | 69.59% | 70.90% | 67.96% | 71.94% | 70.30% | 69.98% | 69.62% |
| Statin Therapy for Patients With Diabetes Statin Adherence 80% (Total) ² | 74.78% | 74.09% | 68.73% | 73.64% | 75.75% | 65.65% | 79.37% | 75.04% | 73.22% | 73.79% |
| Asthma Medication Ratio, Ages 5 to 64 Years ³ | 67.25% | 62.87% | 67.65% | 68.80% | 64.19% | 59.55% | 68.78% | 66.29% | 65.43% | 64.79% |
| Controlling High Blood Pressure, Ages 18 to 85 Years ^{2,3} | 74.45% | 71.05% | 61.56% | 73.24% | 65.94% | 63.50% | 73.97% | 70.34% | 65.22% | 63.43% |
| Persistence of Beta Blocker Treatment After a Heart Attack, 18 Years and Older | 91.16% | 83.69% | 77.68% | 81.53% | 83.33% | 81.48% | 89.74% | 85.31% | 86.55% | 85.91% |
| Hemoglobin A1c Control for Patients With Diabetes - Poor HbA1c Control ^{2,3,5,6} | 32.85% | 29.20% | 35.04% | 33.09% | 36.25% | 34.79% | 29.20% | 32.32% | 36.05% | 38.40% |
| Hemoglobin A1c Control for Patients With Diabetes - HbA1c Control (<8%) ^{2,5} | 56.93% | 60.58% | 54.01% | 56.69% | 52.31% | 55.72% | 63.50% | 58.06% | 54.07% | 51.24% |
| Eye Exams for People with Diabetes ^{2,5} | 58.88% | 67.64% | 50.12% | 54.74% | 49.15% | 51.34% | 64.23% | 57.90% | 55.19% | 53.34% |
| Blood Pressure Control for Patients With Diabetes (<140/90 mm Hg) ^{2,5} | 72.99% | 79.81% | 59.61% | 72.02% | 64.72% | 66.67% | 76.16% | 71.22% | 67.04% | 66.04% |
| Use of Imaging Studies for Low Back Pain, Ages 18 to 64 Years | 72.94% | 73.55% | 79.98% | 72.65% | 79.46% | 75.32% | 76.15% | 75.75% | 76.57% | 77.20% |
| Antidepressant Medication Management (Effective Acute Phase Treatment) | 64.11% | 64.63% | 54.94% | 59.93% | 60.07% | 57.77% | 65.91% | 62.25% | 61.19% | 56.77% |
| Antidepressant Medication Management (Effective Continuation Phase Treatment) | 47.14% | 45.31% | 37.28% | 40.63% | 43.74% | 40.10% | 48.50% | 44.51% | 43.79% | 41.16% |
| Follow-Up Care for Children Prescribed ADHD Medication (Initiation Phase) ² | 45.89% | 42.60% | 55.27% | 44.72% | 33.30% | 42.46% | 53.20% | 45.36% | 41.23% | 47.55% |
| Follow-Up Care for Children Prescribed ADHD Medication (Continuation and Maintenance Phase) ² | 53.37% | 48.01% | 64.02% | 53.66% | 43.46% | 46.62% | 60.43% | 53.25% | 48.93% | 52.84% |
| Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications, Ages 18 to 64 Years | 87.67% | 88.78% | 76.21% | 86.61% | 84.65% | 85.74% | 87.30% | 85.99% | 86.93% | 85.62% |
| Diabetes Monitoring for People With Diabetes and Schizophrenia, Ages 18 to 64 Years | 71.75% | 78.45% | 75.37% | 74.07% | 74.26% | 70.11% | 78.98% | 76.02% | 72.06% | 69.74% |
| Cardiovascular Monitoring for People With Cardiovascular Disease and Schizophrenia, Ages 18 to 64 Years | NA | NA | NA | NA | 90.70% | NA | 78.33% | 81.59% | 76.39% | 72.99% |
| Adherence to Antipsychotic Medications for Individuals With Schizophrenia, Ages 19 to 64 Years | 65.51% | 61.49% | 59.16% | 66.27% | 62.19% | 57.86% | 76.41% | 67.54% | 61.30% | 65.13% |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - Blood Glucose Testing, Total | 79.17% | 83.40% | 54.08% | 80.00% | 72.87% | 71.43% | 79.21% | 77.95% | 76.11% | 69.80% |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - Cholesterol Testing, Total | 69.12% | 75.24% | 60.28% | 72.93% | 64.13% | 61.65% | 68.33% | 69.19% | 66.37% | 60.72% |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - Blood Glucose and Cholesterol Testing, Total | 67.49% | 74.07% | 48.17% | 70.28% | 60.66% | 60.90% | 67.07% | 66.94% | 64.06% | 58.23% |

MY 2022 (RY 2023) HEDIS® Rate Chart

| Measure | Health Plan | | | | | | | MY 2022 HC Weighted Average | MY 2021 HC Weighted Average | MY 2020 HC Weighted Average |
|---|----------------------------------|-----------------------|-----------------------|---------------------------------|----------------|-------------------|--------------|-----------------------------|-----------------------------|-----------------------------|
| | AmeriHealth Caritas Pennsylvania | Geisinger Health Plan | Health Partners Plans | Highmark Wholecare ¹ | Keystone First | United Healthcare | UPMC for You | | | |
| HDO Use of Opioids at High Dosage ^{2,6} | 7.82% | 6.84% | 5.07% | 6.75% | 17.57% | 8.92% | 6.60% | 7.91% | 7.94% | 8.57% |
| Use of Opioids From Multiple Providers - Multiple Pharmacies ^{2,6} | 0.94% | 0.32% | 0.90% | 0.89% | 1.17% | 2.58% | 1.74% | 1.41% | 1.21% | 1.42% |
| Use of Opioids From Multiple Providers - Multiple Prescribers ^{2,6} | 10.48% | 13.42% | 14.73% | 13.14% | 10.98% | 14.29% | 18.06% | 15.71% | 14.05% | 13.64% |
| Use of Opioids From Multiple Providers - Multiple Prescribers and Multiple Pharmacies ^{2,6} | 0.22% | 0.13% | 0.45% | 0.45% | 0.51% | 1.59% | 1.00% | 0.76% | 0.71% | 0.65% |
| Risk of Continued Opioid Use Total >15 Days ^{2,6} | 1.55% | 2.39% | 4.89% | 3.49% | 2.91% | 2.83% | 6.34% | 3.90% | 3.27% | 5.07% |
| Risk of Continued Opioid Use Total >31 Days ^{2,6} | 1.28% | 1.69% | 3.02% | 2.21% | 2.45% | 2.21% | 3.92% | 2.63% | 2.04% | 3.15% |
| Plan All Cause Readmissions Expected Readmission Rate - Total Stays (Ages Total) ⁷ | 9.77% | 9.75% | 9.81% | 9.28% | 9.85% | 10.32% | 9.86% | 9.81% | 9.68% | 9.79% |
| Plan All Cause Readmissions Observed Readmission - Total Stays (Ages Total) ⁷ | 9.05% | 7.63% | 10.49% | 8.54% | 11.16% | 11.54% | 7.52% | 9.42% | 9.37% | 10.02% |
| Plan All Cause Readmission Observed to Expected Readmission Ratio - Total Stays (Ages Total) ^{3,6} | 0.93 | 0.78 | 1.07 | 0.92 | 1.13 | 1.12 | 0.76 | 0.96 | 0.97 | 1.02 |
| Cardiac Rehabilitation - Initiation (18-64) | 6.58% | 3.21% | 1.86% | 1.52% | 1.92% | 2.32% | 2.54% | 2.77% | 2.14% | 2.04% |
| Cardiac Rehabilitation - Engagement1 (18-64) | 8.42% | 5.25% | 1.24% | 0.00% | 2.87% | 4.30% | 4.68% | 3.95% | 2.68% | 2.73% |
| Cardiac Rehabilitation - Engagement2 (18-64) | 6.84% | 5.25% | 1.55% | 0.00% | 2.87% | 4.64% | 4.20% | 3.68% | 2.10% | 2.36% |
| Cardiac Rehabilitation - Achievement (18-64) | 1.84% | 1.17% | 0.00% | 0.00% | 0.38% | 1.99% | 1.95% | 1.18% | 0.37% | 1.08% |
| Cardiac Rehabilitation - Initiation (65+) | NA | NA | NA | NA | NA | NA | 5.71% | 5.71% | NA | NA |
| Cardiac Rehabilitation - Engagement1 (65+) | NA | NA | NA | NA | NA | NA | 12.86% | 12.86% | NA | NA |
| Cardiac Rehabilitation - Engagement2 (65+) | NA | NA | NA | NA | NA | NA | 14.29% | 14.29% | NA | NA |
| Cardiac Rehabilitation - Achievement (65+) | NA | NA | NA | NA | NA | NA | 8.57% | 8.57% | NA | NA |
| Cardiac Rehabilitation - Initiation (Total) | 7.01% | 3.18% | 1.81% | 1.51% | 1.89% | 2.32% | 2.74% | 2.86% | 2.16% | 2.02% |
| Cardiac Rehabilitation - Engagement1 (Total) | 8.83% | 5.20% | 1.21% | 0.00% | 2.83% | 4.30% | 5.21% | 4.17% | 2.74% | 2.73% |
| Cardiac Rehabilitation - Engagement2 (Total) | 6.75% | 5.20% | 1.51% | 0.00% | 2.83% | 4.64% | 4.84% | 3.87% | 2.16% | 2.33% |
| Cardiac Rehabilitation - Achievement (Total) | 1.82% | 1.16% | 0.00% | 0.00% | 0.38% | 1.99% | 2.37% | 1.33% | 0.36% | 1.07% |
| Kidney Health Evaluation for Patient's With Diabetes (18-64) | 43.21% | 55.72% | 44.64% | 41.88% | 43.90% | 42.64% | 45.54% | 45.42% | 41.22% | 38.55% |
| Kidney Health Evaluation for Patient's With Diabetes (65-74) | 53.85% | 70.43% | 53.06% | 53.33% | 51.78% | 47.30% | 53.37% | 53.39% | 50.49% | 45.42% |
| Kidney Health Evaluation for Patient's With Diabetes (75-84) | 53.10% | 78.00% | 51.04% | 55.77% | 50.47% | 53.85% | 48.92% | 51.15% | 49.68% | 40.53% |
| Kidney Health Evaluation for Patient's With Diabetes (Total) | 43.52% | 55.98% | 44.96% | 42.15% | 44.24% | 42.97% | 46.34% | 45.86% | 41.47% | 38.70% |
| Pharmacotherapy for Opioid Use Disorder (16-64) ² | 25.84% | 28.92% | 19.99% | 19.89% | 23.60% | 19.07% | 20.58% | 22.23% | 22.12% | 27.19% |
| Pharmacotherapy for Opioid Use Disorder (65+) ² | NA | NA | NA | NA | NA | NA | 33.82% | 33.82% | NA | NA |
| Pharmacotherapy for Opioid Use Disorder (Total) ² | 25.85% | 28.91% | 20.02% | 19.90% | 23.61% | 19.15% | 20.73% | 22.28% | 22.13% | 27.18% |
| Domain: Access / Availability of Care | | | | | | | | | | |
| Adults' Access to Preventive / Ambulatory Health Services, Ages 20 to 44 Years | 79.24% | 80.03% | 64.90% | 76.00% | 69.32% | 61.94% | 80.46% | 74.29% | 75.53% | 75.15% |
| Adults' Access to Preventive / Ambulatory Health Services, Ages 45 to 64 Years | 85.57% | 86.30% | 77.07% | 83.68% | 79.70% | 71.07% | 87.65% | 83.18% | 82.81% | 82.88% |
| Adults' Access to Preventive / Ambulatory Health Services, Ages 65 Years and Older | 77.79% | 79.43% | 71.95% | 75.15% | 73.18% | 65.16% | 95.23% | 87.25% | 75.74% | 73.30% |
| Adults' Access to Preventive / Ambulatory Health Services, Total | 81.18% | 81.96% | 68.56% | 78.28% | 72.52% | 64.58% | 83.80% | 77.37% | 77.77% | 77.61% |
| Annual Dental Visit, Ages 2 to 20 Years ^{2,3} | 61.28% | 65.41% | 57.66% | 63.92% | 67.17% | 75.31% | 59.68% | 63.23% | 60.50% | 54.23% |
| Prenatal Care in First Trimester ^{2,3} | 90.51% | 88.81% | 87.59% | 88.08% | 87.10% | 89.05% | 90.02% | 88.74% | 88.97% | 88.93% |
| Postpartum Care ^{2,3} | 85.40% | 79.56% | 79.56% | 78.10% | 81.51% | 80.05% | 83.94% | 81.58% | 79.58% | 77.80% |

MY 2022 (RY 2023) HEDIS® Rate Chart

| Measure | Health Plan | | | | | | | MY 2022 HC Weighted Average | MY 2021 HC Weighted Average | MY 2020 HC Weighted Average |
|---|----------------------------------|-----------------------|-----------------------|---------------------------------|----------------|-------------------|--------------|-----------------------------|-----------------------------|-----------------------------|
| | AmeriHealth Caritas Pennsylvania | Geisinger Health Plan | Health Partners Plans | Highmark Wholecare ¹ | Keystone First | United Healthcare | UPMC for You | | | |
| Domain: Utilization and Relative Resource Use | | | | | | | | | | |
| Well-Child Visits in the First 30 Months of Life: Children who Turned 15 Months (6 or more visits) ^{2,3} | 70.48% | 66.02% | 59.31% | 70.39% | 64.18% | 60.07% | 77.79% | 68.12% | 65.30% | 65.19% |
| Well-Child Visits in the First 30 Months of Life: Children who Turned 30 Months (2 or more visits) | 75.44% | 75.34% | 66.26% | 74.55% | 71.02% | 70.66% | 80.57% | 74.02% | 71.63% | 74.61% |
| Child and Adolescent Well-Care Visits, Ages 3 to 11 Years ² | 65.67% | 65.04% | 64.11% | 64.99% | 66.40% | 63.34% | 69.64% | 66.26% | 65.33% | 60.45% |
| Child and Adolescent Well-Care Visits, Ages 12 to 17 Years ² | 59.00% | 57.82% | 58.50% | 58.36% | 60.62% | 57.84% | 63.21% | 59.91% | 59.63% | 54.68% |
| Child and Adolescent Well-Care Visits, Ages 18 to 21 Years ² | 33.82% | 34.33% | 34.03% | 35.62% | 36.88% | 33.53% | 39.00% | 35.90% | 35.56% | 35.04% |
| Child and Adolescent Well-Care Visits, Total ^{2,3} | 58.24% | 56.99% | 56.85% | 57.55% | 59.42% | 55.96% | 62.39% | 58.89% | 58.37% | 54.60% |
| Frequency of Selected Procedures | | | | | | | | | | |
| Bariatric Weight Loss Surgery, Ages 0 to 19 Years - Male ^{7,8} | 0.05 | 0.02 | 0.14 | 0.00 | 0.03 | 0.04 | 0.03 | 0.04 | 0.00 | 0.00 |
| Bariatric Weight Loss Surgery, Ages 0 to 19 Years - Female ^{7,8} | 0.16 | 0.03 | 0.16 | 0.01 | 0.18 | 0.19 | 0.12 | 0.12 | 0.01 | 0.01 |
| Bariatric Weight Loss Surgery, Ages 20 to 44 Years - Male ^{7,8} | 0.99 | 0.61 | 0.61 | 0.50 | 0.45 | 0.75 | 0.58 | 0.64 | 0.05 | 0.03 |
| Bariatric Weight Loss Surgery, Ages 20 to 44 Years - Female ^{7,8} | 7.23 | 3.66 | 4.29 | 3.98 | 3.18 | 5.91 | 3.76 | 4.57 | 0.34 | 0.28 |
| Bariatric Weight Loss Surgery, Ages 45 to 64 Years - Male ^{7,8} | 1.28 | 0.78 | 0.55 | 0.54 | 0.57 | 0.65 | 0.60 | 0.71 | 0.06 | 0.05 |
| Bariatric Weight Loss Surgery, Ages 45 to 64 Years - Female ^{7,8} | 4.76 | 3.03 | 3.26 | 3.22 | 1.93 | 3.20 | 3.20 | 3.23 | 0.24 | 0.23 |
| Tonsillectomy, Ages 0 to 9 Years ^{7,8} | 5.89 | 5.21 | 3.69 | 4.36 | 3.96 | 3.81 | 5.32 | 4.61 | 0.26 | 0.38 |
| Tonsillectomy, Ages 10 to 19 Years ^{7,8} | 2.35 | 2.17 | 1.36 | 2.23 | 1.39 | 1.52 | 2.61 | 1.95 | 0.14 | 0.17 |
| Hysterectomy, Abdominal, Ages 15 to 44 Years ^{7,8} | 0.75 | 0.71 | 0.45 | 0.58 | 0.66 | 0.29 | 0.60 | 0.58 | 0.06 | 0.06 |
| Hysterectomy, Abdominal, Ages 45 to 64 Years ^{7,8} | 1.65 | 1.45 | 1.16 | 1.57 | 2.01 | 0.92 | 1.29 | 1.44 | 0.12 | 0.14 |
| Hysterectomy, Vaginal, Ages 15 to 44 Years ^{7,8} | 0.96 | 0.51 | 0.49 | 0.78 | 0.36 | 0.51 | 0.99 | 0.66 | 0.05 | 0.07 |
| Hysterectomy, Vaginal, Ages 45 to 64 Years ^{7,8} | 1.23 | 0.79 | 1.28 | 1.14 | 1.13 | 0.62 | 0.99 | 1.03 | 0.09 | 0.09 |
| Cholecystectomy, Closed (Laparoscopic), Ages 30 to 64 Years - Male ^{7,8} | 2.39 | 2.76 | 1.15 | 2.05 | 1.22 | 1.71 | 3.33 | 2.09 | 0.20 | 0.21 |
| Cholecystectomy, Closed (Laparoscopic), Ages 15 to 44 Years - Female ^{7,8} | 7.24 | 6.66 | 3.56 | 5.91 | 3.20 | 4.39 | 7.00 | 5.42 | 0.48 | 0.48 |
| Cholecystectomy, Closed (Laparoscopic), Ages 45 to 64 Years - Female ^{7,8} | 5.80 | 5.90 | 2.66 | 6.00 | 3.54 | 4.92 | 5.80 | 4.95 | 0.46 | 0.44 |
| Cholecystectomy, Open, Ages 30 to 64 Years - Male ^{7,8} | 0.30 | 0.24 | 0.17 | 0.30 | 0.26 | 0.31 | 0.24 | 0.26 | 0.02 | 0.02 |
| Cholecystectomy, Open, Ages 15 to 44 Years - Female ^{7,8} | 0.13 | 0.06 | 0.13 | 0.12 | 0.11 | 0.07 | 0.03 | 0.09 | 0.01 | 0.01 |
| Cholecystectomy, Open, Ages 45 to 64 Years - Female ^{7,8} | 0.26 | 0.21 | 0.08 | 0.24 | 0.08 | 0.25 | 0.37 | 0.21 | 0.02 | 0.02 |
| Back Surgery, Ages 20 to 44 Years - Male ^{7,8} | 1.40 | 1.98 | 1.10 | 1.85 | 1.06 | 1.19 | 1.97 | 1.51 | 0.16 | 0.15 |
| Back Surgery, Ages 20 to 44 Years - Female ^{7,8} | 1.69 | 1.70 | 0.68 | 1.54 | 0.92 | 1.34 | 1.94 | 1.40 | 0.13 | 0.13 |
| Back Surgery, Ages 45 to 64 Years - Male ^{7,8} | 6.63 | 6.68 | 2.81 | 5.54 | 3.87 | 4.66 | 7.08 | 5.32 | 0.46 | 0.48 |
| Back Surgery, Ages 45 to 64 Years - Female ^{7,8} | 5.41 | 5.81 | 2.55 | 5.33 | 3.25 | 4.18 | 7.34 | 4.84 | 0.46 | 0.43 |
| Mastectomies, Ages 15 to 44 Years - Female ^{7,8} | 0.76 | 0.58 | 0.85 | 0.53 | 1.46 | 0.80 | 0.76 | 0.82 | 0.07 | 0.06 |
| Mastectomies, Ages 45 to 64 Years - Female ^{7,8} | 1.78 | 1.54 | 1.28 | 1.06 | 1.85 | 1.97 | 1.42 | 1.56 | 0.16 | 0.15 |
| Lumpectomies, Ages 15 to 44 Years - Female ^{7,8} | 1.33 | 1.25 | 1.00 | 0.87 | 1.07 | 0.87 | 1.10 | 1.07 | 0.10 | 0.10 |
| Lumpectomies, Ages 45 to 64 Years - Female ^{7,8} | 4.60 | 2.91 | 2.85 | 2.90 | 3.75 | 3.44 | 3.16 | 3.37 | 0.28 | 0.29 |
| Ambulatory Care, All Ages | | | | | | | | | | |
| Outpatient Visits (Total) ⁹ | 4465.76 | 4200.27 | 2947.98 | 3954.34 | 3125.11 | 3300.22 | 5414.43 | 4036.89 | 325.72 | 307.80 |
| Emergency Department Visits (Total) ^{6,8} | 723.99 | 568.30 | 556.32 | 661.06 | 556.10 | 580.50 | 652.70 | 617.71 | 50.69 | 48.76 |

MY 2022 (RY 2023) HEDIS® Rate Chart

| Measure | Health Plan | | | | | | | MY 2022 HC Weighted Average | MY 2021 HC Weighted Average | MY 2020 HC Weighted Average |
|--|----------------------------------|-----------------------|-----------------------|---------------------------------|----------------|-------------------|--------------|-----------------------------|-----------------------------|-----------------------------|
| | AmeriHealth Caritas Pennsylvania | Geisinger Health Plan | Health Partners Plans | Highmark Wholecare ¹ | Keystone First | United Healthcare | UPMC for You | | | |
| Inpatient Utilization - General Hospital/Acute Care, All Ages | | | | | | | | | | |
| Total Discharges (Total) ^{7,9} | 63.70 | 62.33 | 60.49 | 69.14 | 76.05 | 62.98 | 80.16 | 67.84 | 5.92 | 6.05 |
| Maternity Discharges (Total) ^{7,9} | 26.07 | 22.06 | 22.90 | 24.47 | 25.10 | 20.82 | 22.49 | 23.42 | 2.15 | 2.51 |
| Surgery Discharges (Total) ^{7,9} | 14.51 | 13.76 | 15.07 | 17.23 | 15.13 | 15.48 | 24.04 | 16.46 | 1.43 | 1.40 |
| Medicine Discharges (Total) ^{7,9} | 29.84 | 31.71 | 27.78 | 33.61 | 42.37 | 31.45 | 39.36 | 33.73 | 2.89 | 2.81 |
| Total Average Length of Stay (Total) ^{9,10} | 4.86 | 4.08 | 4.25 | 5.05 | 4.80 | 5.40 | 4.68 | | | |
| Maternity Average Length of Stay (Total) ^{9,10} | 2.75 | 2.48 | 1.98 | 2.76 | 2.81 | 2.75 | 2.51 | | | |
| Surgery Average Length of Stay (Total) ^{9,10} | 7.38 | 6.24 | 8.44 | 8.22 | 8.57 | 9.05 | 6.89 | | | |
| Medicine Average Length of Stay (Total) ^{9,10} | 5.00 | 3.99 | 3.42 | 4.67 | 4.32 | 4.95 | 4.25 | | | |
| Antibiotic Utilization | | | | | | | | | | |
| Antibiotic Utilization for Respiratory Conditions - (3m-17) ⁴ | 18.67 | 23.81 | 11.17 | 18.71 | 12.61 | 17.73 | 23.66 | 18.84 | | |
| Antibiotic Utilization for Respiratory Conditions - (18-64) ⁴ | 16.41 | 21.55 | 10.31 | 15.05 | 11.37 | 14.80 | 19.97 | 16.22 | | |
| Antibiotic Utilization for Respiratory Conditions - (65+) ⁴ | 8.23 | 12.92 | 10.09 | 9.05 | 9.24 | 9.30 | 17.38 | 14.13 | | |
| Antibiotic Utilization for Respiratory Conditions - (Total) ⁴ | 17.64 | 22.78 | 10.73 | 17.10 | 12.06 | 16.33 | 21.89 | 17.62 | | |
| Domain: Health Plan Descriptive Information | | | | | | | | | | |
| HealthChoices Enrollment | 15.81% | 11.01% | 12.41% | 12.72% | 19.45% | 4.46% | 24.14% | | | |
| Domain: Measures Reported Using Electronic Clinical Data Systems | | | | | | | | | | |
| Childhood Immunization Status - DTaP - E ⁴ | 69.97% | 74.65% | 66.45% | 68.17% | 72.08% | 69.70% | 72.91% | 70.84% | | |
| Childhood Immunization Status - IPV - E ⁴ | 84.35% | 89.71% | 80.47% | 84.98% | 85.98% | 84.13% | 87.43% | 85.50% | | |
| Childhood Immunization Status - MMR - E ⁴ | 85.67% | 87.52% | 85.15% | 85.93% | 87.06% | 85.40% | 87.02% | 86.39% | | |
| Childhood Immunization Status - Hib - E ⁴ | 82.13% | 84.81% | 83.14% | 82.62% | 86.24% | 82.92% | 86.10% | 84.37% | | |
| Childhood Immunization Status - Hepatitis B - E ⁴ | 81.49% | 89.55% | 82.71% | 84.26% | 85.62% | 84.85% | 86.48% | 85.01% | | |
| Childhood Immunization Status - VZV - E ⁴ | 85.25% | 86.75% | 84.92% | 85.60% | 87.08% | 84.99% | 86.78% | 86.10% | | |
| Childhood Immunization Status - Pneumococcal Conjugate - E ⁴ | 72.64% | 77.77% | 68.08% | 71.05% | 72.91% | 71.83% | 76.45% | 73.21% | | |
| Childhood Immunization Status - Hepatitis A - E ⁴ | 80.86% | 82.66% | 84.13% | 81.38% | 85.44% | 82.27% | 83.99% | 83.25% | | |
| Childhood Immunization Status - Rotavirus - E ⁴ | 66.31% | 71.50% | 62.04% | 66.71% | 70.23% | 68.20% | 72.35% | 68.67% | | |
| Childhood Immunization Status - Influenza - E ⁴ | 42.57% | 39.80% | 46.43% | 38.98% | 52.09% | 48.58% | 41.84% | 44.67% | | |
| Childhood Immunization Status - Combo 3 - E ⁴ | 61.76% | 68.83% | 60.83% | 60.42% | 66.31% | 63.80% | 66.30% | 64.28% | | |
| Childhood Immunization Status - Combo 7 - E ⁴ | 52.22% | 58.22% | 51.46% | 50.85% | 57.97% | 54.86% | 57.78% | 55.18% | | |
| Childhood Immunization Status - Combo 10 - E ⁴ | 29.00% | 30.48% | 31.68% | 26.25% | 39.48% | 35.96% | 31.62% | 32.49% | | |
| Immunizations for Adolescents - Meningococcal - E ⁴ | 85.51% | 85.44% | 80.57% | 85.20% | 86.79% | 83.48% | 86.20% | 85.13% | | |
| Immunizations for Adolescents - Tdap - E ⁴ | 85.74% | 85.93% | 81.89% | 85.69% | 87.41% | 84.32% | 86.47% | 85.69% | | |
| Immunizations for Adolescents - HPV - E ⁴ | 37.46% | 32.98% | 42.82% | 34.42% | 45.47% | 37.58% | 35.92% | 38.72% | | |
| Immunizations for Adolescents - Combination 1 - E ⁴ | 84.64% | 84.43% | 79.34% | 84.33% | 85.92% | 82.64% | 85.32% | 84.20% | | |
| Immunizations for Adolescents - Combination 2 - E ⁴ | 36.73% | 32.39% | 41.41% | 33.81% | 44.64% | 37.09% | 35.44% | 37.99% | | |
| Breast Cancer Screening - E ¹¹ | 58.02% | 57.26% | 54.42% | 50.50% | 53.30% | 48.33% | 56.25% | 54.95% | 50.77% | |
| Colorectal Cancer Screening (Total) - E ⁴ | 39.54% | 41.54% | 32.81% | 38.37% | 33.34% | 26.44% | 43.94% | 38.40% | | |

MY 2022 (RY 2023) HEDIS® Rate Chart

| Measure | Health Plan | | | | | | | MY 2022 HC Weighted Average | MY 2021 HC Weighted Average | MY 2020 HC Weighted Average |
|--|----------------------------------|-----------------------|-----------------------|---------------------------------|----------------|-------------------|--------------|-----------------------------|-----------------------------|-----------------------------|
| | AmeriHealth Caritas Pennsylvania | Geisinger Health Plan | Health Partners Plans | Highmark Wholecare ¹ | Keystone First | United Healthcare | UPMC for You | | | |
| Follow-Up Care for Children Prescribed ADHD Medication - Initiation Phase - E ¹¹ | 45.89% | 42.60% | 55.27% | 44.69% | 33.26% | 42.46% | 53.20% | 45.35% | 41.10% | |
| Follow-Up Care for Children Prescribed ADHD Medication -Continuation Phase - E ¹¹ | 53.37% | 48.01% | 64.02% | 53.56% | 43.33% | 46.62% | 60.43% | 53.21% | 48.54% | |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - Blood Glucose Testing (Total) - E ⁴ | 79.17% | 83.40% | 52.96% | 80.00% | 72.87% | 71.43% | 79.21% | 77.91% | | |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - Cholesterol Testing - (Total) - E ⁴ | 69.12% | 75.24% | 60.28% | 72.93% | 64.13% | 61.65% | 68.33% | 69.19% | | |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics - Blood Glucose and Cholesterol Testing (Total) - E ⁴ | 67.49% | 74.07% | 47.32% | 70.28% | 60.66% | 60.90% | 67.07% | 66.91% | | |
| Depression Screening and Follow-Up for Adolescents and Adults - Depression Screening (12-17) - E ¹¹ | 1.56% | 14.17% | 2.22% | 0.01% | 1.34% | 0.09% | 1.81% | 2.83% | 2.11% | |
| Depression Screening and Follow-Up for Adolescents and Adults - Depression Screening (18-64) - E ⁴ | 3.18% | 11.21% | 4.06% | 0.14% | 4.09% | 0.54% | 2.57% | 3.75% | | |
| Depression Screening and Follow-Up for Adolescents and Adults - Depression Screening (65+) - E ⁴ | 4.23% | 8.26% | 1.73% | 0.00% | 1.72% | 0.00% | 2.58% | 2.46% | | |
| Depression Screening and Follow-Up for Adolescents and Adults - Depression Screening (Total) - E ⁴ | 2.81% | 11.85% | 3.70% | 0.11% | 3.42% | 0.46% | 2.43% | 3.54% | | |
| Depression Screening and Follow-Up for Adolescents and Adults - Follow-up on Positive Screen (12-17) - E ¹¹ | 67.80% | 75.92% | 41.77% | NA | 73.53% | NA | 89.67% | 59.60% | 59.86% | |
| Depression Screening and Follow-Up for Adolescents and Adults - Follow-up on Positive Screen (18-64) - E ⁴ | 62.03% | 68.78% | 46.92% | NA | 61.90% | NA | 75.58% | 61.46% | | |
| Depression Screening and Follow-Up for Adolescents and Adults - Follow-up on Positive Screen (65+) - E ⁴ | NA | NA | NA | NA | NA | NA | 58.33% | 40.67% | | |
| Depression Screening and Follow-Up for Adolescents and Adults - Follow-up on Positive Screen (Total) - E ⁴ | 63.41% | 70.87% | 46.03% | NA | 63.18% | NA | 77.16% | 62.42% | | |
| Adult Immunization Status - Influenza - E ¹¹ | 17.81% | 17.80% | 12.92% | 15.85% | 17.75% | 15.46% | 17.92% | 16.80% | 18.46% | |
| Adult Immunization Status - TdTap - E ¹¹ | 48.99% | 55.18% | 39.69% | 48.87% | 39.22% | 48.62% | 46.95% | 45.91% | 42.67% | |
| Adult Immunization Status - Zoster - E ¹¹ | 13.16% | 16.17% | 5.15% | 10.48% | 10.97% | 10.87% | 12.19% | 11.38% | 8.37% | |
| Prenatal Immunization Status - Influenza - E | 31.30% | 29.47% | 33.49% | 27.30% | 32.47% | 30.11% | 27.75% | 30.27% | 33.63% | 39.73% |
| Prenatal Immunization Status - Tdap - E | 71.50% | 71.63% | 67.53% | 67.02% | 66.77% | 62.88% | 69.01% | 68.32% | 67.59% | 67.98% |
| Prenatal Immunization Status - Combination - E | 28.26% | 26.90% | 29.41% | 24.23% | 28.28% | 26.04% | 24.67% | 26.83% | 29.26% | 34.25% |
| Prenatal Depression Screening and Follow-Up - Depression Screening - E ¹¹ | 26.50% | 50.84% | 39.65% | 3.27% | 44.38% | 19.82% | 29.92% | 31.61% | 26.41% | |
| Prenatal Depression Screening - Follow-Up on Positive Prenatal Depression Screening - E ¹¹ | 49.21% | 63.31% | 43.27% | NA | 45.61% | 50.00% | 56.88% | 50.77% | 41.33% | |
| Postpartum Depression Screening and Follow-Up - Depression Screening - E ¹¹ | 27.58% | 39.35% | 44.91% | 4.03% | 39.96% | 3.13% | 38.57% | 30.49% | 21.79% | |
| Postpartum Depression Screening - Follow-Up on Positive Postpartum Depression Screening - E ¹¹ | 54.17% | 60.53% | 46.67% | NA | 48.06% | NA | 67.76% | 59.75% | 48.25% | |

* Rates in red font are less than the weighted average for the measure and may appear equal to weighted averages due to rounding, unless they are inverted measures.

** NA signifies denominator is less than 30.

*** It is DHS's practice to report all first-year performance measures for informational purposes, which may not align with NCQA's practice for reporting first-year performance measures.

¹ Effective 01/01/2022, this information is issued on behalf of Highmark Wholecare, coverage by Gateway Health Plan, which is an independent licensee of the Blue Cross Blue Shield Association. Highmark Wholecare serves a Medicaid plan to Blue Shield members in 13 counties in central Pennsylvania, as well as, to Blue Cross Blue Shield members in 27 counties in western Pennsylvania.

² NCQA advises to trend MY 2020 weighted averages with caution. For measure specific details, please reference the memo issued by NCQA: https://www.ncqa.org/wp-content/uploads/2021/02/20210226_HEDIS_MY_2020_Measure_Review_Memo.pdf.

³ Measures highlighted in yellow are MCO Pay-for-Performance (P4P) measures.

⁴ MY 2022 is first year of public reporting by DHS.

⁵ Effective MY 2022, NCQA revised the CDC submeasures to standalone measures without making substantive changes to the measure specifications.

⁶ A lower rate indicates better performance for these measures.

⁷ These measures are not assigned a HC weighted average per NCQA's HEDIS guidance that higher or lower service counts do not indicate better or worse performance impacting the ability to meaningfully interpret averages. Instead, these measures are assigned a PA DHS Mean.

⁸ Beginning MY 2022, the formula for this measure changed to N/D x 12000 to produce member year data. Previously, for MY 2020 and MY 2021, the formula was N/D x 1000. Trending should be done with caution.

⁹ Beginning MY 2022, the discharge formula for this measure changed to Discharges/Member Months x 12000 to produce member year data. Previously, for MY 2020 and MY 2021, the formula was Discharges/Member Months x 1000. Trending should be done with caution.

¹⁰ These measures are not based on a population of members in the denominator and NCQA does not view higher or lower service counts as an indication of better or worse performance impacting the ability to meaningfully interpret averages. These measures are also already presented for each MCO as an average.

¹¹ MY 2021 is first year of public reporting by DHS.